

3. Spinale FG, Mukherjee R, Zavadzka JA, Koval CN, Bouges S, Stroud RE, et al. Cardiac restricted overexpression of membrane type-1 matrix metalloproteinase causes adverse myocardial remodeling following myocardial infarction. *J Biol Chem.* 2010;285:30316-27.
4. d'Ortho MP, Will H, Atkinson S, Butler G, Messent A, Gavrilovic J, et al. Membrane-type matrix metalloproteinases 1 and 2 exhibit broad-spectrum proteolytic capacities comparable to many matrix metalloproteinases. *Eur J Biochem.* 1997; 250:751-7.
5. Knauper V, Will H, Lopez-Otin C, Smith B, Atkinson SJ, Stanton H, et al. Cellular mechanisms for human procollagenase-3 (MMP-13) activation. Evidence that MT1-MMP (MMP-14) and gelatinase A (MMP-2) are able to generate active enzyme. *J Biol Chem.* 1996;271:17124-31.
6. Spruill LS, Lowry AS, Stroud RE, Squires CE, Mains IM, Flack EC, et al. Membrane-type-1 matrix metalloproteinase transcription and translation in myocardial fibroblasts from patients with normal left ventricular function and from patients with cardiomyopathy. *Am J Physiol Cell Physiol.* 2007;293:C1362-73.
7. van Rooij E, Olson EN. MicroRNAs: powerful new regulators of heart disease and provocative therapeutic targets. *J Clin Invest.* 2007;117:2369-76.
8. van Rooij E, Sutherland LB, Thatcher JE, DiMaio JM, Naseem RH, Marshall WS, et al. Dysregulation of microRNAs after myocardial infarction reveals a role of miR-29 in cardiac fibrosis. *Proc Natl Acad Sci U S A.* 2008;105: 13027-32.
9. Duisters RF, Tijssen AJ, Schroein B, Leenders JJ, Lentink V, van der Made I, et al. miR-133 and miR-30 regulate connective tissue growth factor: implications for a role of microRNAs in myocardial matrix remodeling. *Circ Res.* 2009;104: 170-8, 6p following 8.
10. Matkovich SJ, Wang W, Tu Y, Eschenbacher WH, Dorn LE, Condorelli G, et al. MicroRNA-133a protects against myocardial fibrosis and modulates electrical repolarization without affecting hypertrophy in pressure-overloaded adult hearts. *Circ Res.* 2010;106:166-75.
11. Friedman RC, Farh KK, Burge CB, Bartel DP. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res.* 2009;19:92-105.
12. Grimson A, Farh KK, Johnston WK, Garrett-Engle P, Lim LP, Bartel DP. MicroRNA targeting specificity in mammals: determinants beyond seed pairing. *Mol Cell.* 2007;27:91-105.
13. Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell.* 2005;120:15-20.
14. Tatti O, Vehviläinen P, Lehti K, Keski-Oja J. MT1-MMP releases latent TGF-beta1 from endothelial cell extracellular matrix via proteolytic processing of LTBP-1. *Exp Cell Res.* 2008;314:2501-14.
15. Roberts AB, Sporn MB, Assoian RK, Smith JM, Roche NS, Wakefield LM, et al. Transforming growth factor type beta: rapid induction of fibrosis and angiogenesis in vivo and stimulation of collagen formation in vitro. *Proc Natl Acad Sci U S A.* 1986;83:4167-71.
16. Wrana JL, Attisano L, Carcamo J, Zentella A, Doody J, Laiho M, et al. TGF beta signals through a heteromeric protein kinase receptor complex. *Cell.* 1992;71: 1003-14.
17. Karsdal MA, Larsen L, Engsig MT, Lou H, Ferreras M, Lochter A, et al. Matrix metalloproteinase-dependent activation of latent transforming growth factor-beta controls the conversion of osteoblasts into osteocytes by blocking osteoblast apoptosis. *J Biol Chem.* 2002;277:44061-7.
18. Rodriguez D, Morrison CJ, Overall CM. Matrix metalloproteinases: what do they not do? New substrates and biological roles identified by murine models and proteomics. *Biochim Biophys Acta.* 2010;1803:39-54.
19. Rehmsmeier M, Steffen P, Hochsmann M, Giegerich R. Fast and effective prediction of microRNA/target duplexes. *RNA.* 2004;10:1507-17.
20. Flack EC, Lindsey ML, Squires CE, Kaplan BS, Stroud RE, Clark LL, et al. Alterations in cultured myocardial fibroblast function following the development of left ventricular failure. *J Mol Cell Cardiol.* 2006;40:474-83.
21. Wang K, Zhang S, Weber J, Baxter D, Galas DJ. Export of microRNAs and microRNA-protective protein by mammalian cells. *Nucleic Acids Res.* 2010; 38:7248-59.
22. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods.* 2001; 25:402-8.
23. Chapman RE, Spinale FG. Extracellular protease activation and unraveling of the myocardial interstitium: critical steps toward clinical applications. *Am J Physiol Heart Circ Physiol.* 2004;286:H1-10.
24. Spinale FG. Myocardial matrix remodeling and the matrix metalloproteinases: influence on cardiac form and function. *Physiol Rev.* 2007;87:1285-342.
25. Guo C, Piacentini L. Type I collagen-induced MMP-2 activation coincides with up-regulation of membrane type 1-matrix metalloproteinase and TIMP-2 in cardiac fibroblasts. *J Biol Chem.* 2003;278:46699-708.
26. Kim JI, Cordova AC, Hirayama Y, Madri JA, Sumpio BE. Differential effects of shear stress and cyclic strain on Sp1 phosphorylation by protein kinase C-zeta modulates membrane type 1-matrix metalloproteinase in endothelial cells. *Endothelium.* 2008;15:33-42.
27. Ruddy JM, Jones JA, Stroud RE, Mukherjee R, Spinale FG, Ikonomidis JS. Differential effects of mechanical and biological stimuli on matrix metalloproteinase promoter activation in the thoracic aorta. *Circulation.* 2009;120(11 Suppl):S262-8.
28. Zile MR, Baicu CF, Stroud RE, Van Laer A, Arroyo J, Mukherjee R, et al. Pressure overload-dependent membrane type 1-matrix metalloproteinase induction: relationship to LV remodeling and fibrosis. *Am J Physiol Heart Circ Physiol.* 2012;302:H1429-37.
29. Fontana L, Chen Y, Prijatelj P, Sakai T, Fassler R, Sakai LY, et al. Fibronectin is required for integrin alphavbeta6-mediated activation of latent TGF-beta complexes containing LTBP-1. *FASEB J.* 2005;19:1798-808.
30. Ye Y, Perez-Polo JR, Qian J, Birnbaum Y. The role of microRNA in modulating myocardial ischemia-reperfusion injury. *Physiol Genomics.* 2011;43:534-42.
31. Bostjancic E, Zidar N, Stajer D, Glavac D. MicroRNAs miR-1, miR-133a, miR-133b and miR-208 are dysregulated in human myocardial infarction. *Cardiology.* 2010;115:163-9.
32. Ikeda S, Pu WT. Expression and function of microRNAs in heart disease. *Curr Drug Targets.* 2010;11:913-25.
33. Jones JA, Stroud RE, O'Quinn EC, Black LE, Barth JL, Eleftheriades JA, et al. Selective microRNA suppression in human thoracic aneurysms: relationship of miR-29a to aortic size and proteolytic induction. *Circ Cardiovasc Genet.* 2011; 4:605-13.
34. Tang Y, Zheng J, Sun Y, Wu Z, Liu Z, Huang G. MicroRNA-1 regulates cardiomyocyte apoptosis by targeting Bcl-2. *Int Heart J.* 2009;50:377-87.
35. Slezak S, Jin P, Caruccio L, Ren J, Bennett M, Zia N, et al. Gene and microRNA analysis of neutrophils from patients with polycythemia vera and essential thrombocytosis: down-regulation of micro RNA-1 and -133a. *J Transl Med.* 2009;7:39.
36. He B, Xiao J, Ren AJ, Zhang YF, Zhang H, Chen M, et al. Role of miR-1 and miR-133a in myocardial ischemic postconditioning. *J Biomed Sci.* 2011;18:22.
37. Dorn GW 2nd. MicroRNAs: redefining mechanisms in cardiac disease. *J Cardiovasc Pharmacol.* 2010;56:589-95.
38. Spinale FG, Koval CN, Deschamps AM, Stroud RE, Ikonomidis JS. Dynamic changes in matrix metalloproteinase activity within the human myocardial interstitium during myocardial arrest and reperfusion. *Circulation.* 2008;118(14 Suppl):S16-23.
39. Barallobre-Barreiro J, Didangelos A, Schoendube FA, Drozdov I, Yin X, Fernandez-Caggiano M, et al. Proteomics analysis of cardiac extracellular matrix remodeling in a porcine model of ischemia/reperfusion injury. *Circulation.* 2012; 125:789-802.
40. Jayawardena TM, Egemnazarov B, Finch EA, Zhang L, Payne JA, Pandya K, et al. MicroRNA-mediated in vitro and in vivo direct reprogramming of cardiac fibroblasts to cardiomyocytes. *Circ Res.* 2012;110:1465-73.
41. Brown RD, Ambler SK, Mitchell MD, Long CS. The cardiac fibroblast: therapeutic target in myocardial remodeling and failure. *Annu Rev Pharmacol Toxicol.* 2005;45:657-87.
42. Frangogiannis NG, Smith CW, Entman ML. The inflammatory response in myocardial infarction. *Cardiovasc Res.* 2002;53:31-47.
43. Porter KE, Turner NA. Cardiac fibroblasts: at the heart of myocardial remodeling. *Pharmacol Ther.* 2009;123:255-78.

Discussion

Dr Paul W. M. Fedak (Calgary, Alberta, Canada). There is clear clinical evidence that even with early and successful reperfusion after coronary occlusion, patients will undergo progressive cardiac remodeling that can lead to heart failure. The key molecular and cellular changes that occur after reperfusion are not well known. It gives us a window of opportunity where we could potentially treat patients early during or after reperfusion to prevent maladaptive signals from happening, but we need to understand them better. Your work is important because it leads us in

that direction to better understand mechanistically what is happening with those molecular and cellular changes.

What is unique and important is that you have been able to separate out region- and time-specific changes in a preclinical relevant animal model. What is really important is that you are able to separate ECM interstitial signaling from the cellular changes. It is difficult to present that complexity of data in such a short period of time, and you did a great job. I have a couple of questions for you.

You have designed your experiment to look at mechanistic data and tried to tease out time- and region-specific changes and correlate those events. I note with your data you are trying to correlate MT1-MMP activity and LTB processing, but in the remote area during reperfusion you observed the highest levels of the MMP actively, but you didn't observe the similar changes in the LTB processing. Does that refute your hypothesis?

Dr Eckhouse. I believe the differences seen in MT1-MMP and LTBP-1 activity with microdialysis actually support our hypothesis.

First, in examining MT1-MMP activity with microdialysis during the I/R protocol, there are several possible reasons that could lead to this increased MT1-MMP activity in the remote region. Previous studies from other laboratories have determined that a phosphorylation event occurring at the C-terminus of MT1-MMP can lead to a substrate specificity and directed localization within a cell membrane. From these findings, our research group hypothesizes that post-translational modifications of MT1-MMP in vivo are promoting increased cleavage of the MT1-MMP fluorogenic peptide in a region-dependent manner.

An important question to consider is how are the MT1-MMP and LTBP-1 substrates different? The peptide sequences of each substrate are different. The MT1-MMP fluorogenic peptide is a synthetic amino acid sequence that is hydrolyzed specifically by MT1-MMP in vitro and in vivo. Previous validation studies confirmed that the amino acid sequence is specific for MT1-MMP over other MMPs. The LTBP-1 fluorogenic peptide represents an endogenous substrate sequence that is hydrolyzed by MT1-MMP. Previous in vivo studies have determined that with overexpression of MT1-MMP, there is a decrease in full-length LTBP-1 and an increase in the low-molecular-weight split products of LTBP-1 after MT1-MMP-mediated hydrolysis. Therefore, MT1-MMP differentially processes the 2 synthetic peptides.

The cleavage of the LTBP-1 fluorogenic peptide was specific to the I/R region, which is supported by the increase in the activation of the TGF- β profibrotic pathway as measured by pSmad2 in the I/R region when compared with the remote region. Although not in the presentation, gene expression of collagen 1A1 by quantitative PCR was increased, which further suggests increased acute activation of the TGF- β profibrotic pathway within the I/R region that was not measured within the remote region, as demonstrated by no change in collagen 1A1 gene expression and a small increase in pSmad2 protein abundance when compared with the referent control.

Dr Fedak. Can you speculate what the triggers could be here? Could wall stress be the key trigger, and if so, can you think of anything we can do from a surgical point of view early after reperfusion?

Dr Eckhouse. Wall stress has been demonstrated to be an important mechanism in activating MT1-MMP. We have seen this

in a recent murine model of pressure overload hypertrophy from our laboratory published by Dr Zile and in in vitro aortic wall stress studies.

The alteration in stress-strain patterns due to an acute I/R event is causing an increase in MT1-MMP activation. Therefore, decreasing wall stress due to I/R remains important.

For a future surgical technique, it will be important to improve stress-strain patterns. Although this may be more relevant to a chronic infarction, improving the thickness of the affected myocardium within the I/R injury region by injections of polymers or other method is currently under investigation. However, there is a lot of area for creativity.

Dr Fedak. Thank you.

Dr Joseph D. Schmoker (*Burlington, Vt.*). To follow up on Paul's question about regional issues, there is a mechanism for uncoupling of TGF- β via mechanical transduction through the integrin network, I believe. I was wondering if you came across that mechanism and if that could serve as an explanation for why you are seeing this only in the region of I/R?

Dr Eckhouse. I haven't come across that mechanism specifically within this study, but one of my predecessors in the laboratory, Dr Ruddy, looked at the TGF- β pathway within in vitro studies of aortic wall stress. We are currently investigating increased TGF- β activation secondary to increased aortic wall stress in models of aortic aneurysms. However, I haven't looked at that within the I/R myocardial model, but that is definitely an area to look at.

Dr Schmoker. My other question relates to miR-133a, which has been used as a biomarker in clinical trials. Could it be that the interstitial levels you are measuring are simply secondary to cell injury or death secondary to reperfusion? Have you looked at that? In other words, do you have histology on your I/R region?

Dr Eckhouse. I don't have histology data to present. We did cell culture studies using primary porcine fibroblasts undergoing simulated in vitro I/R, and what we found when we measured miR-133a within the conditioned media is consistent with our in vivo evidence. When it comes to cell death, previous studies using that same I/R model did not show increased cardiomyocyte death, and we didn't have increased fibroblast cell death with our simulated hypoxia reoxygenation model.

Concurrently, with potential cell death within the I/R region, I would have expected miR-133a extracellularly to increase more with apoptosis and cell death and spillage of the intracellular miR-133a into the interstitial space. However, with I/R within that I/R region, we actually see a decrease during the ischemic time of miR-133a.

Dr Schmoker. Right, but I think it is in the reperfusion episode where you are really going to get the cell injury or death.

One last point. Cardiac myocytes are enriched with miR-133a, but I don't know the relative abundance of that miR-133a in fibroblasts. So I am wondering why you didn't use cardiomyocytes in your cell culture.

Dr Eckhouse. That is a study we are currently doing. We haven't completed that study. We have been able to show a correlation between MT1-MMP levels and miR-133a in myocardial fibroblasts. So we are able to demonstrate that miR-133a is within other cells besides the cardiomyocytes, but we are specifically looking at the cardiomyocytes now.

Dr Stephen McKellar (*Rochester, Minn*). Elegant study and model. Good presentation. You presented your I/R model here, but have you looked at this or similar phenomena in an infarction model with sampling in the infarct, border zones, and then distantly? This is an elegant way of studying that region. Is that something you could at least speculate on what you might find in that model?

Dr Eckhouse. A previous study evaluated MT1-MMP activity within an infarction model with a subsequent I/R event 2 weeks after the infarction. MT1-MMP activity increased with infarction and then subsequently increased even more after a subsequent I/R event 2 weeks later. However, in regard to miR-133a, we haven't looked at the long-term effects of I/R on miR-133a levels, but it is definitely something we are interested in.

Dr Thomas K. Waddell (*Toronto, Ontario, Canada*). My question is along the lines of Dr Schmoker's question about the release of miR-133. You mentioned that the exportation was identified as a mechanism of regulation, but on the 2 slides you showed, it was clear that the Y axis was different. So when you measured by PCR the levels of the miR-133 in the interstitium versus the total, what was the scale of that difference? I would put it to you that if the

total is a million, then what you are measuring in the interstitium is 10. It probably doesn't reflect regulation. What I would like you to think about is perhaps it is a mechanism of communication—the exportation of miR-133 allows that to be diffused outward. If that is the case, you might actually see increasing penumbra of decreasing miR-133 levels. So you may want to look at your remote area a little bit as proximal and distant remote.

Dr Eckhouse. We didn't look at proximal and distant remote regions of the myocardium. That is an interesting point and definitely a suggestion we will take to the laboratory. Our data with our miR exportation seem to fit nicely with our acute TGF- β activation. In the remote region, we saw a decrease in the interstitial miR-133a levels, and it remained decreased with reperfusion, which in comparison with total myocardial pools leads us to believe that miR-133a remains elevated intracellularly. Thus, miR-133a can inhibit or attenuate the TGF- β profibrotic pathway within the remote region. Within the I/R region, the extracellular increase in miR-133a may be associated with a intracellular decrease of miR-133a, which may then allow activation of the TGF- β profibrotic signaling pathways and promote fibrosis over time.